



INDUSTRIAL
MATHEMATICS
INSTITUTE

2004:02

Non-trivial t-intersection in the
function lattice

P. Erdős, A. Seress and L.A.
Székely

IMI
Preprint Series

Department of Mathematics
University of South Carolina

Non-trivial t -intersection in the function lattice *

Péter L. Erdős

A. Rényi Institute of Mathematics
Hungarian Academy of Sciences,
Budapest, P.O. Box 127, H-1364 Hungary
elp@renyi.hu

Ákos Seress

Department of Mathematics
The Ohio State University
Columbus, OH 43210
akos@math.ohio-state.edu

László A. Székely

Department of Mathematics
University of South Carolina
Columbia, SC 29208
szekely@math.sc.edu

Abstract

The function lattice, or generalized Boolean algebra, is the set of ℓ -tuples with the i th coordinate an integer between 0 and a bound n_i . Two ℓ -tuples t -intersect if they have at least t common nonzero coordinates. We prove a Hilton–Milner type theorem for systems of t -intersecting ℓ -tuples.

*The work of the first author was partially supported by Hungarian NSF grants T37846, T34702. The work of the second author was partially supported by the NSF grant CCR-0097995. The work of the third author was partially supported by the NSF grants 007 2187 and 030 2307.

1 Introduction

Let t, ℓ , and $n_1 \leq n_2 \leq \dots \leq n_\ell$ be positive integers. Denote by $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ the set of all ℓ -tuples

$$\{\mathbf{k} = (k_1, \dots, k_\ell) : 0 \leq k_i \leq n_i \quad 1 \leq i \leq \ell\}.$$

The *support* of an ℓ -tuple \mathbf{k} is the set of the non-zero coordinates: $\text{supp}(\mathbf{k}) = \{i : k_i \neq 0\}$. We can define a partial ordering on $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ by $\mathbf{j} \leq \mathbf{k}$ if $\text{supp}(\mathbf{j}) \subset \text{supp}(\mathbf{k})$ and for all $i \in \text{supp}(\mathbf{j})$ we have $j_i = k_i$. This partially ordered set is called the *function lattice* (see for example [5]). Another frequently used name is *generalized Boolean algebra*, because the case $n_1 = n_\ell = 1$, i.e., when all n_i are equal to 1, is just the case of (characteristic vectors of) set systems on an ℓ -element underlying set.

We say that two ℓ -tuples \mathbf{j} and \mathbf{k} are *t-intersecting* if there are at least t different integers $i \in \text{supp}(\mathbf{j}) \cap \text{supp}(\mathbf{k})$ such that $j_i = k_i$, or, with other words, if there is an ℓ -tuple \mathbf{t} with support of size t such that $\mathbf{t} \leq \mathbf{k}$ and $\mathbf{t} \leq \mathbf{j}$. Denote by $m_t(n_1, \dots, n_\ell)$ the maximum cardinality of t -intersecting ℓ -tuples in $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ and by $M_t(n_1, \dots, n_\ell)$ the set of all t -intersecting families with this cardinality. The problems to determine the value $m_t(n_1, \dots, n_\ell)$ and to describe the structures of the families in $M_t(n_1, \dots, n_\ell)$, have a very long and notable history even in the case $n_\ell > 1$, and this is the case we are concentrating on in this note.

We start with the history of the case $t = 1$. C. Berge (1974, [4]) determined $m_t(n_1, \dots, n_\ell)$ and $M_t(n_1, \dots, n_\ell)$ when all ℓ -tuples have ℓ -element supports. Different proofs of Berge's result were given by Hsieh (1975, [19]), by Livingston (1979, [21]) in the case when $n_1 = n_\ell$. The first result for set systems with uniform support size different from ℓ , but with $n_1 = n_\ell$, is due to Frankl (published in 1983, [9]). Moreover, Engel (1984, [10]) handled the case with $n_1 = n_\ell$, when the supports of the ℓ -tuples are arbitrary. In fact, Engel proved a Bollobás-type inequality (in the spirit of [8]) for the set of intersecting ℓ -tuples; a simpler proof of this last result is due to P. L. Erdős, U. Faigle and W. Kern (1992, [12]). In 2001 C. Bey gave a complete solution to the $t = 1$ case, for arbitrary n_i 's and any uniform support size (2001, [6]), using his general weighted intersection theorem. This case shows interesting connections to the complete intersection theorem of R. Ahlswede and L. Khachatrian ([2]).

For arbitrary values of t , the first result is due to D. Kleitman (1966 [20]) in the case when $n_1 = n_\ell = 2$, and all supports are of size ℓ . Then P. Frankl

and Z. Füredi handled the case $t \geq 15$, all supports are of size ℓ , and $n_1 = n_\ell$ (1980, [14]), using Frankl’s version of the Erdős-Ko-Rado theorem (see [11]). Later A. Moon generalized this result for cross t -intersecting families (1982, [22]). The paper by Deza and Frankl (1983, [9]) also contains the solution for the case when all supports are of the same size k and $n_1 = n_\ell$, for ℓ large enough as a function of k and t . H-D. Gronau proved the first result for t -intersecting families with ℓ -element supports in the case of non-equal n_i ’s (1983, [16]). R. Ahlswede and L. Khachatrian (1998, [3]), and independently P. Frankl and N. Tokushige (1998, [15]), solved the t -intersecting problem for arbitrary t for ℓ -tuples with full support, applying Ahlswede and Khachatrian’s seminal complete intersection theorem for set systems (1997, [2]). Finally C. Bey (1999, [5]) determined all parameters ℓ, k, t, n , for which “fixing t coordinates” yields the solution to the intersection problem.

All these results can be summarized in the following structural way: under some conditions for the parameter values, the (often unique) optimal t -intersecting family consists of all ℓ -tuples that are greater or equal than a fixed ℓ -tuple \mathbf{t} with support size t . In the literature such set systems are called *trivially* t -intersecting families. As it is well known in the theory of t -intersecting set systems, there is a long-standing effort to solve the *nontrivial* t -intersection problem: what is the size and the structure of the maximum t -intersecting families where the total intersection of the sets has less than t elements. The first such result is due to A. J. H. Hilton and E. C. Milner (1967, [18]). The complete solution is again due to R. Ahlswede and L. Khachatrian (1996, [1]).

As far as these authors are aware, the only t -intersection result known for the function lattice $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ is due to C. Bey and K. Engel (2000, [7]) [Example 10, 11 and Lemma 18]: this is the complete solution to the non-trivial t -intersection problem in the case of equal n_i ’s.

The goal of this paper is to prove a more general non-trivial t -intersection result for the subset of the function lattice $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ consisting of ℓ -tuples with a fixed size k of the support, for some parameter values $t < k < \ell$ and $n_1 \leq n_2 \leq \dots \leq n_\ell$. The result is based on a Hilton–Milner type theorem for poset series, proved by the authors (2000, [13]). The proof of this latter uses the so-called *kernel method*, introduced by A. Hajnal and B. Rothschild (1973, [17]), therefore all of our results are valid only from a threshold for the parameters. We note that, perhaps surprisingly, the

application of [13] is *not* for the natural partial order of $\mathbb{F}_\ell(n_1, \dots, n_\ell)$. We shall investigate families of intersecting chains in the natural partial order of $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ in a forthcoming paper. Of course, a direct application of the kernel method may yield similar results, but citing [13] saves a lot of work. We admit that the methods of [7] are likely to allow generalization to the case of different n_i 's.

In Section 2 we recall the necessary details from [13], while in Section 3 we reformulate the t -intersection problem of the function lattice and apply for it the method described in Section 2.

2 Non-trivial t -intersection results for posets

A t -chain \mathcal{L} in a poset P is a strict chain of elements $\mathcal{L} = (x_1 < x_2 < \dots < x_t)$. For a given t -chain $\mathcal{L} = (x_1 < x_2 < \dots < x_t)$, let $\mathcal{T}_{P,k}(x_1, x_2, \dots, x_t)$ denote the set of k -chains in P which contain \mathcal{L} as a subset. Define $T_{P,k}(x_1, x_2, \dots, x_t) = |\mathcal{T}_{P,k}(x_1, x_2, \dots, x_t)|$. Sometimes we write T instead of $T_{P,k}$, when it does not cause ambiguity. Also define $r_t(P, k) = \max T_{P,k}(x_1, x_2, \dots, x_t)$, where the maximum is taken for t -chains $x_1 < x_2 < \dots < x_t$ in P . It follows from the definition that

$$r_i(P, k) \geq r_{i+1}(P, k). \quad (1)$$

For a t -chain $\mathcal{X} \subset P$ and $y \notin \mathcal{X}$, let $T(\mathcal{X}, y)$ denote the number of k -chains which contain \mathcal{X} and y . For a t -chain \mathcal{X} and a k -chain \mathcal{L} in P , such that $|\mathcal{X} \cup \mathcal{L}| = k + 1$, let $y_{\mathcal{L}}^* \in \mathcal{L} \setminus \mathcal{X}$ such that $T(\mathcal{X}, y_{\mathcal{L}}^*)$ minimize $T(\mathcal{X}, y)$ for the elements $y \in \mathcal{L} \setminus \mathcal{X}$, and set

$$\tau(\mathcal{X}, \mathcal{L}) = \sum_{y \in \mathcal{L} \setminus \mathcal{X}, y \neq y_{\mathcal{L}}^*} T(\mathcal{X}, y). \quad (2)$$

Also define

$$M_\tau(P, k) = \max_{\mathcal{X}, \mathcal{L}} \tau(\mathcal{X}, \mathcal{L}), \quad (3)$$

and

$$M_\tau^*(P, k) = \max_{\substack{\mathcal{X}, \mathcal{L}: \\ \tau(\mathcal{X}, \mathcal{L}) = M_\tau(P, k)}} T(\mathcal{X}, y_{\mathcal{L}}^*). \quad (4)$$

Now the following Hilton-Milner type theorem holds:

Theorem 1 For fixed $1 \leq t < k$, and a sequence of posets P_n , let us be given a maximum sized family \mathcal{F}_n of non-trivially t -intersecting k -chains in P_n . Assume further that

$$\lim_{n \rightarrow \infty} r_{t+2}(P_n, k) / M_\tau^*(P_n, k) = 0. \quad (5)$$

Then, for n sufficiently large, \mathcal{F}_n has one of the following two descriptions:

- (i) there exists a t -chain \mathcal{X} and a $(k+1-t)$ -chain \mathcal{Y} , such that $\mathcal{X} \cap \mathcal{Y} = \emptyset$; and \mathcal{F}_n is the following set of k -chains:

$$\begin{aligned} \mathcal{F}(\mathcal{X}, \mathcal{Y}) &= \{\mathcal{L} : \mathcal{X} \subseteq \mathcal{L} \text{ and } \mathcal{L} \cap \mathcal{Y} \neq \emptyset\} \cup \\ &\cup \{\mathcal{L} : \mathcal{Y} \subseteq \mathcal{L} \text{ and } |\mathcal{L} \cap \mathcal{X}| = t - 1\}, \end{aligned} \quad (6)$$

where the second set of chains is non-empty;

- (ii) there exists a $(t+2)$ -chain \mathcal{Z} , and \mathcal{F}_n is the following set of k -chains:

$$\mathcal{F}(\mathcal{Z}) = \{\mathcal{L} : |\mathcal{L} \cap \mathcal{Z}| \geq t + 1\}, \quad (7)$$

and $|\bigcap_{\mathcal{L} \in \mathcal{F}_n} \mathcal{L} \cap \mathcal{Z}| \leq t - 1$.

3 New results

Let $t < k < \ell$ and $n_1 \leq \dots \leq n_\ell$ be positive integers. We define two families $\mathcal{F}_1(t, k; n_1, \dots, n_\ell)$ and $\mathcal{F}_2(t, k; n_1, \dots, n_\ell)$ of non-trivially t -intersecting families in $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ with support size k as follows.

- (i) Let j_1, j_2, \dots, j_{k+1} be integers satisfying $1 \leq j_i \leq n_i$ for $i \in [1, k+1]$. We define $\mathcal{F}_1(t, k; n_1, \dots, n_\ell)$ as the set of ℓ -tuples $\mathbf{k} = (k_1, \dots, k_\ell)$ with support size k which belong to the set

$$\begin{aligned} &\{\mathbf{k} : k_i = j_i \text{ for all } i \in [1, t] \text{ and for at least one } i \in [t+1, k+1]\} \cup \\ &\{\mathbf{k} : k_i = j_i \text{ for all } i \in [t+1, k+1] \text{ and for } t-1 \text{ values } i \in [1, t]\}. \end{aligned} \quad (8)$$

- (ii) Let j_1, j_2, \dots, j_{t+2} be integers satisfying $1 \leq j_i \leq n_i$ for $i \in [1, t+2]$. We define $\mathcal{F}_2(t, k; n_1, \dots, n_\ell)$ as the set of ℓ -tuples $\mathbf{k} = (k_1, \dots, k_\ell)$ with support size k which belong to the set

$$\{\mathbf{k} : k_i = j_i \text{ for at least } t+1 \text{ values } i \in [1, t+2]\}. \quad (9)$$

Note that $|\mathcal{F}_1(t, k; n_1, \dots, n_\ell)|$ and $|\mathcal{F}_2(t, k; n_1, \dots, n_\ell)|$ do not depend on the particular choices of the j_i . Our goal is to give sufficient conditions for the parameter values $t, k, \ell, n_1, \dots, n_\ell$ which ensure that either \mathcal{F}_1 or \mathcal{F}_2 is of maximum size among the non-trivially t -intersecting families of ℓ -tuples with support size k .

Given $n_1 \leq \dots \leq n_\ell$, we define a partially ordered set $(\mathcal{P}(n_1, \dots, n_\ell), \prec)$ as follows. The underlying set is $\mathcal{P}(n_1, \dots, n_\ell) := \{(i, j) : 1 \leq i \leq \ell, 1 \leq j \leq n_i\}$, and $(i_1, j_1) \prec (i_2, j_2)$ if and only if $i_1 < i_2$. The map $\mathbf{k} = (k_1, \dots, k_\ell) \mapsto \{(i, k_i) \in \mathcal{P}(n_1, \dots, n_\ell) : k_i \neq 0\}$ is obviously a bijection between $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ and the chains in the poset $(\mathcal{P}(n_1, \dots, n_\ell), \prec)$, and ℓ -tuples with support size k are mapped to k -chains. Therefore, t -intersecting families of ℓ -tuples in $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ with support size k correspond to t -intersecting k -chains in $(\mathcal{P}(n_1, \dots, n_\ell), \prec)$. For a subset $\mathcal{Y} \subseteq \mathcal{P}(n_1, \dots, n_\ell)$, we define the *support* of \mathcal{Y} as the set of first coordinates of the elements of \mathcal{Y} ; namely, $\text{supp}(\mathcal{Y}) = \{i \leq \ell : \exists j \leq n_i (i, j) \in \mathcal{Y}\}$. We start with the determination of the quantities r_{t+2} , M_τ , and M_τ^* defined in Section 2. Note that for any m -chain \mathcal{L} in $\mathcal{P} = (\mathcal{P}(n_1, \dots, n_\ell), \prec)$, we have

$$T_{\mathcal{P},k}(\mathcal{L}) = \sum_{\substack{AC[1,\ell] \setminus \text{supp}(\mathcal{L}): \\ |A|=k-m}} \prod_{i \in A} n_i. \quad (10)$$

Proposition 2 *Let $t < k < \ell$, let $\mathcal{P} = (\mathcal{P}(n_1, \dots, n_\ell), \prec)$ and let \mathcal{L} be an m -chain in \mathcal{P} . Suppose that $(i, k_i) \in \mathcal{L}$ and $j \notin \text{supp}(\mathcal{L})$ with $j < i$, and let $\mathcal{L}^* = (\mathcal{L} \setminus \{(i, k_i)\}) \cup \{(j, k_j)\}$ for some $k_j \leq n_j$. Then $T_{\mathcal{P},k}(\mathcal{L}^*) \geq T_{\mathcal{P},k}(\mathcal{L})$, with equality if and only if $n_j = n_{j+1} = \dots = n_i$.*

Proof. We obtain $T_{\mathcal{P},k}(\mathcal{L}^*)$ from $T_{\mathcal{P},k}(\mathcal{L})$ by replacing each occurrence of n_j by n_i in the sum in (10). Hence the inequalities $n_j \leq n_{j+1} \leq \dots \leq n_i$ imply both assertions of the proposition. \square

Let $\sigma_i(x_1, x_2, \dots, x_m)$ denote the i^{th} elementary symmetric polynomial in variables x_1, x_2, \dots, x_m . We define $\sigma_0(x_1, x_2, \dots, x_m) = 1$.

Lemma 3 *Let $t < k < \ell$ and let $\mathcal{P} = (\mathcal{P}(n_1, \dots, n_\ell), \prec)$. Then*

$$r_{t+2}(\mathcal{P}, k) = \sum_{\substack{AC[t+3,\ell]: \\ |A|=k-t-2}} \prod_{i \in A} n_i = \sigma_{k-t-2}(n_{t+3}, \dots, n_\ell). \quad (11)$$

Proof. Proposition 2 implies that for $(t + 2)$ -chains \mathcal{L} in \mathcal{P} , the quantity $T_{\mathcal{P},k}(\mathcal{L})$ is maximized when $\text{supp}(\mathcal{L}) = [1, t + 2]$. \square

Lemma 4 *Let $t < k < \ell$ and let $\mathcal{P} = (\mathcal{P}(n_1, \dots, n_\ell), \prec)$. Then for any t -chain \mathcal{X} and k -chain \mathcal{L} in \mathcal{P} with $|\mathcal{X} \cup \mathcal{L}| = k + 1$, we have $M_\tau(\mathcal{P}, k) = \tau(\mathcal{X}, \mathcal{L})$ if and only if the multiset relations $\{n_i : i \in \text{supp}(\mathcal{X})\} = \{n_i : 1 \leq i \leq t\}$ and $\{n_i : i \in \text{supp}(\mathcal{L})\} \supseteq \{n_i : t + 1 \leq i \leq k\}$ hold.*

Proof. We first note that the condition $|\mathcal{X} \cup \mathcal{L}| = k + 1$ implies that \mathcal{X} and \mathcal{L} have $t - 1$ common elements and $|\mathcal{L} \setminus \mathcal{X}| = k - t + 1$. Moreover, since $\tau(\mathcal{X}, \mathcal{L})$ is the sum of only $k - t$ values $T(\mathcal{X}, y)$ with $y \in \mathcal{L} \setminus \mathcal{X}$, it is possible that for a fixed t -chain \mathcal{X} , $\tau(\mathcal{X}, \mathcal{L})$ is maximized for some \mathcal{L} even though $T(\mathcal{X}, y) = 0$ for some $y \in \mathcal{L} \setminus \mathcal{X}$.

For a fixed t -chain \mathcal{X} , Proposition 2 implies that $\tau(\mathcal{X}, \mathcal{L})$ is maximized for a k -chain \mathcal{L} whose support contains the $k - t$ smallest elements of $[1, \ell] \setminus \text{supp}(\mathcal{X})$. Moreover, another application of Proposition 2 shows that if \mathcal{X}' is obtained by replacing an element $(i_1, j_1) \in \mathcal{X}$ with some (i_2, j_2) satisfying $i_2 < i_1$ and i_2 the smallest number not in $\text{supp}(\mathcal{X})$ then $\tau(\mathcal{X}', \mathcal{L}') \geq \tau(\mathcal{X}, \mathcal{L})$ for an optimal \mathcal{L}' constructed in the way described in the previous sentence. Hence $M_\tau(\mathcal{P}, k) = \tau(\mathcal{X}, \mathcal{L})$ for \mathcal{X}, \mathcal{L} with $\text{supp}(\mathcal{X}) = [1, t]$ and $\text{supp}(\mathcal{L}) \supseteq [t + 1, k]$. Finally, Proposition 2 also implies that if $\text{supp}(\mathcal{X}') \neq [1, t]$ or $\text{supp}(\mathcal{L}') \not\supseteq [t + 1, k]$ then $\tau(\mathcal{X}', \mathcal{L}') < M_\tau(\mathcal{P}, k)$, unless the condition about the multiset of n_i values described in the statement of the lemma holds. \square

Lemma 5 *Let $t < k < \ell$ and let $\mathcal{P} = (\mathcal{P}(n_1, \dots, n_\ell), \prec)$. Then*

$$M_\tau^*(\mathcal{P}, k) = \sum_{\substack{A \subset [t+1, \ell] \setminus \{k+1\}: \\ |A| = k-t-1}} \prod_{i \in A} n_i = \sigma_{k-t-1}(n_{t+1}, \dots, \widehat{n_{k+1}}, \dots, n_\ell). \quad (12)$$

Proof. Let \mathcal{X} be a t -chain and \mathcal{L} be a k -chain with $|\mathcal{X} \cup \mathcal{L}| = k + 1$ and $\tau(\mathcal{X}, \mathcal{L}) = M_\tau(\mathcal{P}, k)$. Then, by Lemma 4, we have the multiset relations $\{n_i : i \in \text{supp}(\mathcal{X})\} = \{n_i : 1 \leq i \leq t\}$ and $\{n_i : i \in \text{supp}(\mathcal{L})\} \supseteq \{n_i : t + 1 \leq i \leq k\}$. Also, we have $k \leq |\text{supp}(\mathcal{X} \cup \mathcal{L})| \leq k + 1$. If $|\text{supp}(\mathcal{X} \cup \mathcal{L})| = k$ then there exists $y_{\mathcal{L}}^* = (i, k_i) \in \mathcal{L} \setminus \mathcal{X}$ with $i \in \text{supp}(\mathcal{X})$ and so $T(\mathcal{X}, y_{\mathcal{L}}^*) = 0$. If $|\text{supp}(\mathcal{X} \cup \mathcal{L})| = k + 1$ then Proposition 2 implies that $T(\mathcal{X}, y)$ is minimized in $\mathcal{L} \setminus \mathcal{X}$ for the $y_{\mathcal{L}}^* = (i, k_i) \in \mathcal{L} \setminus \mathcal{X}$ with $i = \max \text{supp}(\mathcal{L} \setminus \mathcal{X})$ and, in order to maximize $T(\mathcal{X}, y_{\mathcal{L}}^*)$, we have to choose $\max \text{supp}(\mathcal{L} \setminus \mathcal{X})$ as small as possible.

Combining these observations, we obtain that $\max T(\mathcal{X}, y_{\mathcal{L}}^*)$ is achieved in the case $\text{supp}(\mathcal{X}) = [1, t]$, $\text{supp}(\mathcal{L} \setminus \mathcal{X}) = [t+1, k+1]$, and $\text{supp}(y_{\mathcal{L}}^*) = \{k+1\}$, leading to (12). \square

The following two lemmas will be useful at the comparison of r_{t+2} and M_{τ}^* .

Lemma 6 *Let t, k, ℓ satisfy $k \geq t+2$ and $\ell \geq 2k - t - 1$, and let $\mathcal{P} = (\mathcal{P}(n_1, \dots, n_{\ell}), \prec)$. Then*

$$r_{t+2}(\mathcal{P}, k) \leq \left(1 + \frac{k-t-2}{\ell-2k+t+2}\right) \sigma_{k-t-2}(n_{t+1}, \dots, \widehat{n_{k+1}}, \dots, n_{\ell}).$$

Proof. On one hand, if $A \subseteq [t+1, \ell]$ satisfies $|A| = k-t-2$ and $k+1 \in A$ then

$$\prod_{i \in A} n_i \leq \frac{\sum_{s \in [k+2, \ell] \setminus A} n_s}{(\ell-k-1) - (k-t-3)} \prod_{i \in A \setminus \{k+1\}} n_i.$$

On the other hand, any $(k-t-2)$ -element subset B of $[t+1, \ell] \setminus \{k+1\}$ can be obtained at most $k-t-2$ ways by replacing $k+1$ by an element $j \geq k+2$ of B . Hence Lemma 3 implies

$$\begin{aligned} r_{t+2}(\mathcal{P}, k) &= \sigma_{k-t-2}(n_{t+3}, \dots, n_{\ell}) \leq \sigma_{k-t-2}(n_{t+1}, \dots, n_{\ell}) \leq \\ &\left(1 + \frac{k-t-2}{\ell-2k+t+2}\right) \sigma_{k-t-2}(n_{t+1}, \dots, \widehat{n_{k+1}}, \dots, n_{\ell}). \end{aligned}$$

\square

Lemma 7 *Let t, k, ℓ satisfy $k \geq t+2$ and let $\mathcal{P} = (\mathcal{P}(n_1, \dots, n_{\ell}), \prec)$. Then*

$$M_{\tau}^*(\mathcal{P}, k) \geq n_{t+1} \frac{\ell-k+1}{k-t-1} \sigma_{k-t-2}(n_{t+1}, \dots, \widehat{n_{k+1}}, \dots, n_{\ell}). \quad (13)$$

Proof. Using that any $(k-t-2)$ -element subset B of $[t+1, \ell] \setminus \{k+1\}$ can be obtained $(\ell-t-1) - (k-t-2) = \ell-k+1$ ways by deleting an element different from $k+1$ from a $(k-t-1)$ -element subset of $[t+1, \ell] \setminus \{k+1\}$, we have

$$\begin{aligned} &(k-t-1) \sigma_{k-t-1}(n_{t+1}, \dots, \widehat{n_{k+1}}, \dots, n_{\ell}) = \\ &\sum_{\substack{s=t+1 \\ s \neq k+1}}^{\ell} n_s \sigma_{k-t-2}(n_{t+1}, \dots, \widehat{n_s}, \dots, \widehat{n_{k+1}}, \dots, n_{\ell}) \geq \\ &n_{t+1} (\ell-k+1) \sigma_{k-t-2}(n_{t+1}, \dots, \widehat{n_{k+1}}, \dots, n_{\ell}). \end{aligned}$$

Hence Lemma 5 implies (13). \square

Lemma 8 *Let $t < k < \ell$ and let $\mathcal{P} = (\mathcal{P}(n_1, \dots, n_\ell), \prec)$. If \mathcal{X} is a t -chain and \mathcal{Y} is a $k+1-t$ -chain with $\mathcal{X} \cap \mathcal{Y} = \emptyset$ then $|\mathcal{F}(\mathcal{X}, \mathcal{Y})| \leq |\mathcal{F}_1(t, k; n_1, \dots, n_\ell)|$ for the families of chains defined in (6) and (8), respectively.*

Proof. First note that $|\text{supp}(\mathcal{X}) \cap \text{supp}(\mathcal{Y})| \leq 1$, because otherwise there is no k -chain containing \mathcal{Y} and $t-1$ elements of \mathcal{X} as required in (6). If $|\text{supp}(\mathcal{X}) \cap \text{supp}(\mathcal{Y})| = 1$, say $(i, f_i) \in \mathcal{X}$ and $(i, g_i) \in \mathcal{Y}$ for some $f_i \neq g_i$, then there exists exactly one k -chain in $\mathcal{F}(\mathcal{X}, \mathcal{Y})$ which contains (i, g_i) , namely, $(\mathcal{Y} \cup \mathcal{X}) \setminus \{(i, f_i)\}$. Hence, if we define $\mathcal{Y}_1 = (\mathcal{Y} \setminus \{(i, g_i)\}) \cup \{(j, 1)\}$ for some $j \notin \text{supp}(\mathcal{X} \cup \mathcal{Y})$ then $|\mathcal{F}(\mathcal{X}, \mathcal{Y})| \leq |\mathcal{F}(\mathcal{X}, \mathcal{Y}_1)|$, because $\mathcal{F}(\mathcal{X}, \mathcal{Y}_1)$ contains all but one chain from $\mathcal{F}(\mathcal{X}, \mathcal{Y})$ and it contains t chains not in $\mathcal{F}(\mathcal{X}, \mathcal{Y})$ (the chains obtained by deleting an element of \mathcal{X} from $\mathcal{X} \cup \mathcal{Y}$). Therefore, it is enough to prove that $|\mathcal{F}(\mathcal{X}, \mathcal{Y})| \leq |\mathcal{F}_1(t, k; n_1, \dots, n_\ell)|$ for chains \mathcal{X}, \mathcal{Y} with $\text{supp}(\mathcal{X}) \cap \text{supp}(\mathcal{Y}) = \emptyset$.

Suppose now that $\text{supp}(\mathcal{X}) \cap \text{supp}(\mathcal{Y}) = \emptyset$. There are exactly t chains in $\mathcal{F}(\mathcal{X}, \mathcal{Y})$ containing \mathcal{Y} and there are t chains in $\mathcal{F}_1(t, k; n_1, \dots, n_\ell)$ with support containing $[t+1, k+1]$; hence it is enough to show that for the set of chains

$$\mathcal{F}^*(\mathcal{X}, \mathcal{Y}) = \{\mathcal{L} : \mathcal{X} \subseteq \mathcal{L} \text{ and } \mathcal{L} \cap \mathcal{Y} \neq \emptyset\}$$

and

$$\mathcal{F}_1^*(t, k; n_1, \dots, n_\ell) = \{\mathcal{L} \in \mathcal{F}_1(t, k; n_1, \dots, n_\ell) : \text{supp}(\mathcal{L}) \supseteq [1, t]\}$$

we have $|\mathcal{F}^*(\mathcal{X}, \mathcal{Y})| \leq |\mathcal{F}_1^*(t, k; n_1, \dots, n_\ell)|$. If $\text{supp}(\mathcal{X}) \neq [1, t]$ then we define a new set of chains by the following shifting operation. Let $i_1 \in [1, t]$ be the smallest number not in $\text{supp}(\mathcal{X})$ and let $i_2 \in \text{supp}(\mathcal{X})$ with $i_2 > i_1$, say $(i_2, k_{i_2}) \in \mathcal{X}$. For a k -chain $\mathcal{L} \in \mathcal{F}^*(\mathcal{X}, \mathcal{Y})$, let

$$f(\mathcal{L}) = \begin{cases} (\mathcal{L} \setminus \{(i_2, k_{i_2})\}) \cup \{(i_1, 1)\} & \text{if } i_1 \notin \text{supp}(\mathcal{L}), \\ (\mathcal{L} \setminus \{(i_1, k_{i_1}), (i_2, k_{i_2})\}) \cup \{(i_1, 1), (i_2, k_{i_1})\} & \text{if } (i_1, k_{i_1}) \in \mathcal{L} \text{ for some } k_{i_1} \leq n_{i_1}. \end{cases} \quad (14)$$

Moreover, define $\mathcal{X}' = (\mathcal{X} \setminus \{(i_2, k_{i_2})\}) \cup \{(i_1, 1)\}$ and

$$\mathcal{Y}' = \begin{cases} \mathcal{Y} & \text{if } i_1 \notin \text{supp}(\mathcal{Y}), \\ (\mathcal{Y} \setminus \{(i_1, k_{i_1})\}) \cup \{(i_2, k_{i_1})\} & \text{if } (i_1, k_{i_1}) \in \mathcal{Y} \text{ for some } k_{i_1} \leq n_{i_1}. \end{cases} \quad (15)$$

Then it is clear that f is an injection from $\mathcal{F}^*(\mathcal{X}, \mathcal{Y})$ into $\mathcal{F}^*(\mathcal{X}', \mathcal{Y}')$, and so $|\mathcal{F}^*(\mathcal{X}, \mathcal{Y})| \leq |\mathcal{F}^*(\mathcal{X}', \mathcal{Y}')|$ and $|\mathcal{F}(\mathcal{X}, \mathcal{Y})| \leq |\mathcal{F}(\mathcal{X}', \mathcal{Y}')|$. Repeating this procedure, we arrive to some t -chain \mathcal{X}'' and $(k+1-t)$ -chain \mathcal{Y}'' such that $|\mathcal{F}^*(\mathcal{X}, \mathcal{Y})| \leq |\mathcal{F}^*(\mathcal{X}'', \mathcal{Y}'')|$ and $\text{supp}(\mathcal{X}'') = [1, t]$ and $\text{supp}(\mathcal{X}'') \cap \text{supp}(\mathcal{Y}'') = \emptyset$. It is enough to show that $|\mathcal{F}^*(\mathcal{X}'', \mathcal{Y}'')| \leq |\mathcal{F}_1^*(t, k; n_1, \dots, n_\ell)|$.

If $\text{supp}(\mathcal{Y}'') \neq [t+1, k+1]$ then let $i_1 \in [t+1, k+1]$ be the smallest number not in $\text{supp}(\mathcal{Y}'')$ and let $i_2 \in \text{supp}(\mathcal{Y}'')$ with $i_2 > i_1$, say $(i_2, k_{i_2}) \in \mathcal{Y}''$. By renumbering the i_2 th coordinate, we may assume that $k_{i_2} \leq n_{i_1}$. We apply the following modification of the shifting operation described in the previous paragraph. For a k -chain $\mathcal{L} \in \mathcal{F}^*(\mathcal{X}'', \mathcal{Y}'')$, let

$$g(\mathcal{L}) = \begin{cases} (\mathcal{L} \setminus \{(i_2, j_2)\}) \cup \{(i_1, j_2)\} & \text{if } i_1 \notin \text{supp}(\mathcal{L}) \text{ and } (i_2, j_2) \in \mathcal{L} \text{ with } j_2 \leq n_1, \\ (\mathcal{L} \setminus \{(i_1, j_1)\}) \cup \{(i_2, j_1)\} & \text{if } i_2 \notin \text{supp}(\mathcal{L}) \text{ and } (i_1, j_1) \in \mathcal{L}, \\ (\mathcal{L} \setminus \{(i_1, j_1), (i_2, j_2)\}) \cup \{(i_1, j_2), (i_2, j_1)\} & \text{if } (i_1, j_1), (i_2, j_2) \in \mathcal{L} \text{ and } j_2 \leq n_1, \\ \mathcal{L} & \text{otherwise.} \end{cases} \quad (16)$$

Moreover, define $\mathcal{Y}''' = (\mathcal{Y}'' \setminus \{(i_2, k_{i_2})\}) \cup \{(i_1, k_{i_2})\}$. Then g is an injection from $\mathcal{F}^*(\mathcal{X}'', \mathcal{Y}'')$ into $\mathcal{F}^*(\mathcal{X}''', \mathcal{Y}''')$, and so $|\mathcal{F}^*(\mathcal{X}'', \mathcal{Y}'')| \leq |\mathcal{F}^*(\mathcal{X}''', \mathcal{Y}''')|$ and $|\mathcal{F}(\mathcal{X}'', \mathcal{Y}'')| \leq |\mathcal{F}(\mathcal{X}''', \mathcal{Y}''')|$. Repeating this procedure, we arrive to a member of the family $\mathcal{F}_1^*(t, k; n_1, \dots, n_\ell)$. \square

Lemma 9 *Let $t < k < \ell$ and let $\mathcal{P} = (\mathcal{P}(n_1, \dots, n_\ell), \prec)$. If \mathcal{Z} is a $(t+2)$ -chain then $|\mathcal{F}(\mathcal{Z})| \leq |\mathcal{F}_2(t, k; n_1, \dots, n_\ell)|$ for the families of chains defined in (7) and (9), respectively.*

Proof. Given $\mathcal{F}(\mathcal{Z})$, if $\text{supp}(\mathcal{Z}) \neq [1, t+2]$ then we can apply the shifting procedure described in (16), not decreasing the size of $\mathcal{F}(\mathcal{Z})$, and eventually arriving to a set of chains in the family $\mathcal{F}_2(t, k; n_1, \dots, n_\ell)$. \square

Lemma 10 *For \mathcal{F}_1 and \mathcal{F}_2 from (8) and (9),*

$$\begin{aligned} |\mathcal{F}_1| &= \sigma_{k-t}(n_{t+1}, \dots, n_\ell) - \sigma_{k-t}(n_{t+1}-1, \dots, n_{k+1}-1, n_{k+2}, \dots, n_\ell) + t \\ |\mathcal{F}_2| &= \sum_{i=1}^{t+2} \sigma_{k-t-1}(n_i, n_{t+3}, \dots, n_\ell) - (t+1)\sigma_{k-t-2}(n_{t+3}, \dots, n_\ell). \end{aligned}$$

Proof. Explanation for $|\mathcal{F}_1|$. The second line of (8) yields the term t , and the cardinality arising from the first line of (8) is obtained as a difference, counting all functions \mathbf{k} with $k_i = j_i$ for all $i \in [1, t]$, and subtracting the number of functions \mathbf{k} with $k_i = j_i$ for all $i \in [1, t]$ that have no $i \in [t + 1, k + 1]$ with $k_i = j_i$.

Explanation for $|\mathcal{F}_2|$. Fix a $(t + 2)$ -chain \mathcal{Z} with support $[1, t + 2]$. For $i \in [1, t + 2]$, the number of k -chains intersecting \mathcal{Z} in coordinates $1, 2, \dots, i - 1, i + 1, \dots, t + 2$ is $\sigma_{k-t-1}(n_i, n_{t+3}, \dots, n_\ell)$. Adding these expressions for all $i \in [1, t + 2]$, the k -chains intersecting \mathcal{Z} in exactly $t + 1$ coordinates are counted once, and the k -chains intersecting \mathcal{Z} in $t + 2$ coordinates are counted $t + 2$ times. The negative term reduces the multiplicity of the latter ones to one. \square

In order to apply Theorem 1, we have to find values of the parameters $t, k, \ell, n_1, \dots, n_\ell$ such that the hypothesis of the theorem is satisfied.

Theorem 11 *Let $t < k < \ell$ be fixed. Then there exists a bound $n(t, k, \ell)$ such that if $n > n(t, k, \ell)$ then for any non-trivially t -intersecting family \mathcal{F} of ℓ -tuples with support k in $\mathbb{F}_\ell(n, \dots, n)$ we have*

$$|\mathcal{F}| \leq \max\{|\mathcal{F}_1(t, k; n, \dots, n)|, |\mathcal{F}_2(t, k; n, \dots, n)|\}.$$

Moreover, if $k > 2t + 1$ then for large enough n we have $|\mathcal{F}_1(t, k; n, \dots, n)| > |\mathcal{F}_2(t, k; n, \dots, n)|$ and if $t + 1 < k \leq 2t + 1$ then for large enough n we have $|\mathcal{F}_1(t, k; n, \dots, n)| < |\mathcal{F}_2(t, k; n, \dots, n)|$.

Proof. Let $\mathcal{P}_n = (\mathcal{P}(n, \dots, n), \prec)$. By Lemmas 3 and 5, we have $r_{t+2}(\mathcal{P}_n, k) = \binom{l-t-2}{k-t-2} n^{k-t-2}$ and $M_\tau^*(\mathcal{P}_n, k) = \binom{l-t-1}{k-t-1} n^{k-t-1}$. Hence

$$\lim_{n \rightarrow \infty} \frac{r_{t+2}(\mathcal{P}_n, k)}{M_\tau^*(\mathcal{P}_n, k)} = \lim_{n \rightarrow \infty} \frac{k-t-1}{l-t-1} \cdot \frac{1}{n} = 0 \quad (17)$$

and so Theorem 1, together with Lemmas 8 and 9, implies that for large enough n one of the maximum sized families of t -intersecting ℓ -tuples with support k in $\mathbb{F}_\ell(n, \dots, n)$ is $\mathcal{F}_1 = \mathcal{F}_1(t, k; n, \dots, n)$ or $\mathcal{F}_2 = \mathcal{F}_2(t, k; n, \dots, n)$.

Our final task is to compare $|\mathcal{F}_1(t, k; n, \dots, n)|$ and $|\mathcal{F}_2(t, k; n, \dots, n)|$. From Lemma 10 we have

$$|\mathcal{F}_1| = t + \binom{l-t}{k-t} n^{k-t} - \sum_{i=0}^{k-t} \binom{k+1-t}{i} \binom{l-k-1}{k-t-i} (n-1)^i n^{k-t-i} \quad (18)$$

and

$$|\mathcal{F}_2| = (t+2) \binom{l-t-1}{k-t-1} n^{k-t-1} - (t+1) \binom{l-t-2}{k-t-2} n^{k-t-2}. \quad (19)$$

Suppose now that $t+2 \leq k$. For fixed t, k, ℓ , as $n \rightarrow \infty$, we expand (18) and (19) as polynomials of n . There is nothing to do with (19), as it is already written in polynomial form. In (18), the coefficient of n^{k-t} in $|\mathcal{F}_1|$ is

$$\binom{l-t}{k-t} - \sum_{i=0}^{k-t} \binom{k+1-t}{i} \binom{l-k-1}{k-t-i} = 0,$$

the coefficient of n^{k-t-1} in $|\mathcal{F}_1|$ is

$$\begin{aligned} & \sum_{i=1}^{k-t} i \binom{k+1-t}{i} \binom{l-k-1}{k-t-i} = \\ & \sum_{i=1}^{k-t} (k+1-t) \binom{k-t}{i-1} \binom{l-k-1}{k-t-i} = (k+1-t) \binom{l-t-1}{k-t-1}, \end{aligned}$$

and similarly the coefficient of n^{k-t-2} in $|\mathcal{F}_1|$ is

$$- \sum_{i=1}^{k-t} \binom{i}{2} \binom{k+1-t}{i} \binom{l-k-1}{k-t-i} = - \frac{(k+1-t)(k-t)}{2} \binom{l-t-2}{k-t-2}.$$

We compare $|\mathcal{F}_1|$ and $|\mathcal{F}_2|$ for large n . The leading term in both is n^{k-t-1} , with coefficients $(k+1-t) \binom{l-t-1}{k-t-1}$ and $(t+2) \binom{l-t-1}{k-t-1}$. Therefore, if $k+1-t > t+2$, i.e. $k > 2t+1$, then for large enough n we have $|\mathcal{F}_1| > |\mathcal{F}_2|$ and if $k < 2t+1$ then for large enough n we have $|\mathcal{F}_1| < |\mathcal{F}_2|$. If $k-t-1 = t+2$, i.e. $k = 2t+1$, then the main terms have equal coefficients. We compare the coefficients of the next term, $n^{k-t-2} = n^{t-1}$ in $|\mathcal{F}_1|$ and $|\mathcal{F}_2|$, which are $-\frac{(t+2)(t+1)}{2} \binom{l-t-2}{t-1}$ and $-(t+1) \binom{l-t-2}{t-1}$, respectively. We have $|\mathcal{F}_1| < |\mathcal{F}_2|$. \square

Theorem 12 *Let $t < k$ be fixed. Then there exists a bound $\ell(t, k)$ such that if $\ell > \ell(t, k)$ then for any non-trivially t -intersecting family \mathcal{F} of ℓ -tuples with support k in $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ we have*

$$|\mathcal{F}| \leq \max\{|\mathcal{F}_1(t, k; n_1, \dots, n_\ell)|, |\mathcal{F}_2(t, k; n_1, \dots, n_\ell)|\}.$$

Proof. Let $\mathcal{P}_\ell = (\mathcal{P}(n_1, \dots, n_\ell), \prec)$. If $k = t + 1$ then $r_{t+2}(\mathcal{P}_\ell, k) = 0$ and $M_\tau^*(\mathcal{P}_\ell, k) > 0$. If $k \geq t + 2$ then by Lemmas 6 and 7, for $\ell \geq 2k - t - 1$ we have

$$\frac{r_{t+2}(\mathcal{P}_\ell, k)}{M_\tau^*(\mathcal{P}_\ell, k)} \leq \left(1 + \frac{k - t - 2}{\ell - 2k + t + 2}\right) \cdot \frac{1}{n_{t+1}} \cdot \frac{k - t - 1}{\ell - k + 1} \quad (20)$$

and therefore

$$\lim_{\ell \rightarrow \infty} \frac{r_{t+2}(\mathcal{P}_\ell, k)}{M_\tau^*(\mathcal{P}_\ell, k)} = 0.$$

So Theorem 1, together with Lemmas 8 and 9, implies that for large enough ℓ one of the maximum sized families of t -intersecting ℓ -tuples with support k in $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ is $\mathcal{F}_1 = \mathcal{F}_1(t, k; n_1, \dots, n_\ell)$ or $\mathcal{F}_2 = \mathcal{F}_2(t, k; n_1, \dots, n_\ell)$. \square

Theorem 13 *Let $t < k < \ell$ be fixed, satisfying $\ell \geq 2k - t - 1$. Then there exists a bound $n(t, k, \ell)$ such that if $n_{t+1} > n(t, k, \ell)$ then for any non-trivially t -intersecting family \mathcal{F} of ℓ -tuples with support k in $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ we have $|\mathcal{F}| \leq \max\{|\mathcal{F}_1(t, k; n_1, \dots, n_\ell)|, |\mathcal{F}_2(t, k; n_1, \dots, n_\ell)|\}$.*

Proof. Let $\mathcal{P}_{n_{t+1}} = (\mathcal{P}(n_1, \dots, n_{t+1}, \dots, n_\ell), \prec)$. If $k = t + 1$ then $r_{t+2}(\mathcal{P}_{n_{t+1}}, k) = 0$ and $M_\tau^*(\mathcal{P}_{n_{t+1}}, k) > 0$. If $k \geq t + 2$ then, analogously to (20) in the proof of Theorem 12,

$$\frac{r_{t+2}(\mathcal{P}_{n_{t+1}}, k)}{M_\tau^*(\mathcal{P}_{n_{t+1}}, k)} \leq \left(1 + \frac{k - t - 2}{\ell - 2k + t + 2}\right) \cdot \frac{1}{n_{t+1}} \cdot \frac{k - t - 1}{\ell - k + 1} \quad (21)$$

and therefore

$$\lim_{n_{t+1} \rightarrow \infty} \frac{r_{t+2}(\mathcal{P}_{n_{t+1}}, k)}{M_\tau^*(\mathcal{P}_{n_{t+1}}, k)} = 0.$$

So Theorem 1, together with Lemmas 8 and 9, implies that for large enough n_{t+1} one of the maximum sized families of t -intersecting ℓ -tuples with support k in $\mathbb{F}_\ell(n_1, \dots, n_\ell)$ is $\mathcal{F}_1 = \mathcal{F}_1(t, k; n_1, \dots, n_\ell)$ or $\mathcal{F}_2 = \mathcal{F}_2(t, k; n_1, \dots, n_\ell)$. \square

Acknowledgement We are indebted to two anonymous referees, who pointed out a minor error in an earlier version of the manuscript.

References

1. R. Ahlswede - L. H. Khachatrian: The complete nontrivial-intersection theorem for systems of finite sets. *J. Comb. Theory (A)* **76** (1996), 121–138.

2. R. Ahlswede - L. H. Khachatrian: The complete intersection theorem for systems of finite sets. *Europ. J. Comb.* **18** (1997), 125–136.
3. R. Ahlswede - L. H. Khachatrian: The diametric theorem in Hamming spaces - Optimal anticodes. *Adv. Appl. Math.* **20** (1998), 429–449.
4. C. Berge: Nombres de coloration de l'hypergraphe h -parti complet, *Hypergraph Seminar*, Proceedings of the First Working Seminar on Hypergraphs, 1972. (Eds. C. Berge and D. Ray-Chaudhuri) Lect. Notes in Math. Vol. 411. Springer-Verlag, Berlin-New York, (1974), 13–20.
5. C. Bey: The Erdős–Ko–Rado bound for the function lattice, *Discrete Appl. Math* **95** (1999), 115–125.
6. C. Bey: An intersection theorem for weighted sets. *Discrete Math.* **235** (2001), 145–150.
7. C. Bey - K. Engel: Old and new results for the weighted t -intersection problem via AK-methods. Althöfer, Ingo (ed.) et al., *Numbers, Information and Complexity*. Dordrecht: Kluwer Academic Publishers. (2000) 45–74.
8. B. Bollobás: Sperner systems consisting of pairs of complementary subsets. *J. Comb. Theory (A)* **15** (1973), 363–366.
9. M. Deza - P. Frankl: Erdős–Ko–Rado theorem – 22 years later. *SIAM J. Alg. Disc. Methods* **4** (1983), 419–431.
10. K. Engel: An Erdős–Ko–Rado Theorem for the subcubes of a cube. *Combinatorica* **4** (1984), 133–140.
11. P. Erdős - C. Ko - R. Rado: Intersection theorems for systems of finite sets. *Quart. J. Math. Oxford Ser. 2* **12** (1961), 313–320.
12. P. L. Erdős - U. Faigle - W. Kern: A group-theoretic setting for some intersecting Sperner families. *Comb. Prob.Comp.* **1** (1992), 323–334.
13. P. L. Erdős - Á. Seress - L. A. Székely: Erdős-Ko-Rado and Hilton-Milner type theorems for intersecting chains in posets. *Combinatorica* **20** (2000), 27–45.

14. P. Frankl - Z. Füredi, The Erdős-Ko-Rado theorem for integer sequences, *SIAM J. Alg. Disc. Methods* **1** (1980), 376–381.
15. P. Frankl - N. Tokushige: The Erdős-Ko-Rado theorem for integer sequence, *Combinatorica* **19** (1999), 55–63.
16. H-D. O. F. Gronau: More on the EKR Theorem for integer sequences. *J. Comb. Theory (A)* **35** (1983), 279–288.
17. A. Hajnal - B. Rothschild: A generalization of the Erdős-Ko-Rado theorem on finite set systems. *J. Comb. Theory (A)* **15** (1973), 359–362.
18. A. J. W. Hilton - E. C. Milner: Some intersection theorems for systems of finite sets. *Quart. J. Math. Oxford (2)* **18** (1967), 369–384.
19. W. N. Hsieh: Systems of finite vector spaces. *Discrete Math.* **12** (1975), 1–16.
20. D. Kleitman: On a combinatorial conjecture of Erdős. *J. Comb. Theory* **1** (1966), 209–214.
21. M. L. Livingston: An ordered version of the Erdős-Ko-Rado theorem. *J. Comb. Theory (A)* **26** (1979), 162–165.
22. A. Moon: An analogue of Erdős-Ko-Rado theorem for Hamming Schemes $H(n, q)$, *J. Comb. Theory (A)* **32** (1982), 386–390.